



An Optimal Design for Inference via Modified Gamma Frailty Distribution

Abdulazeez, Sikiru Adeyinka

Department of Mathematics and Statistics, Kaduna Polytechnic, Kaduna, Nigeria.

E-mail - ysabdul94@gmail.com, Tel : +2348065616611, +23423607591

Abstract

Frailty models play a central role in formulating the effects of covariates on potentially censored failure times and in the joint modelling of incomplete repeated measures and failure times in longitudinal studies. Survival data are often subject to right censoring and to a subsequent loss of information about the effect of explanatory variables. Three frailty models are used to analyze bivariate time-to-event data. All approaches accommodate right censored lifetime data and account for heterogeneity in the study population. A Modified Gamma Frailty Model is compared with two existing Frailty Models. The survival-analysis was performed using the SPSS VERSION 15. The newly derived MGF was analyzed using the Matlab software and R which is more robust when sample size is more than forty. The MGF model performs better than the existing models in the presence of clustering. However the CGF is preferable in the absence of clusters in a given data set.

Keywords: Censorship, Frailty Models, Survival Analysis, Proportional hazard model, Correlated Gamma Frailty Models, Random effects

1. Introduction

Hazard models have become widespread in their use for the analysis of duration time data in many scientific disciplines, including biology and medicine (Cox, 1972; Kalbfleisch and Prentice, 1980), sociology (Petersen, 1998, Vermunt 1996), marketing research (Vilcassim and Jain, 1991; Wedel et al., 1995), (Getachew Y. & Bekele S., 2016) and economics (Kiefer, 1988; Lancaster, 1990). These models overcome the problems of accounting for censored observations of duration and time-varying explanatory variables, that arise in applying standard regression type models to duration data. The basic concept in hazard models is the probability of the occurrence of an event during a certain time interval, say t to $t + \Delta t$, given that it has not occurred before t , specified as:

$$\begin{aligned} \lambda(t|N_i(t-), Z_i(t)) &= \lim_{\Delta t \rightarrow 0} \Pr(t \leq T_{i, N_i(t-)+1} < t + \Delta t | N_i(t-), Z_i(t)) / \Delta t \\ &= \lim_{\Delta t \rightarrow 0} \Pr(\Delta N_i(t) = 1 | N_i(t-), Z_i(t)) / \Delta t \end{aligned}$$

The Cox proportional hazards model (Cox, 1972) is commonly used in the analysis of survival time data. An often unstated assumption of the proportional hazards model and of traditional frailty models (with the exception of those that use the compound Poisson distribution (Aalen, 1988, Aalen, 1992)) is that all individuals will experience the event of interest. However, in some

situation a fraction of individuals is not expected to experience the event of interest; that is, these individuals are not at risk. (Anthony, Isaac & Victor, 2019). The terminology to describe the never-at-risk group varies from field to field, but includes ‘long-term survivors’ or ‘cured’ in epidemiology, ‘non-susceptibles’ in toxicology, ‘stayers’ in finite Markov transition models of occupational mobility, the ‘non-fecundable’ in fertility models, and ‘non-recidivists’ among convicted criminals. In epidemiology and medicine, researchers may be interested in analyzing the occurrence of a disease. Many individuals may never experience that disease; therefore, there exists a fraction in the population that is protected. Cure models are survival models which allow for a cured fraction in the study population.

These models extend the understanding of time-to-event data by allowing for the formulation of more accurate and informative conclusions than previously made. These conclusions would otherwise be unobtainable from an analysis that fails to account for a cured fraction in the population. If a cured component is not present, the analysis reduces to standard approaches of survival analysis. In cure models, the population is divided into two sub-populations so that an individual is either cured with probability $1 - \varphi$, or has a proper survival function $S(t)$, with probability φ . Here, proper survival function means $\lim_{t \rightarrow \infty} S(t) = 0$. Individuals regarded as cured will never experience the event of interest and their survival time will be defined as infinity. Therefore, the hazard and survival functions of cured individuals are set to zero and one, respectively, for all finite values of t . Longini and Halloran (1996) have proposed frailty cure models that extend standard frailty models. The frailty random variable in the former has point mass at zero with probability $1 - \varphi$ while heterogeneity among those experiencing the event of interest is modelled via a continuous distribution with probability φ . Price and Manatunga (2001) gave an excellent introduction to this area and applied leukaemia remission data to different cure, frailty and frailty cure models. They found that frailty models are useful in modelling data with a cured fraction and that the gamma frailty cure model provides a better fit to their remission data compared to the standard cure model. In the next section we describe the existing models and a proposed model, then provide an application of the models to an existing data on occupational exposure tagged – IRANIAN data.

2. Materials and Methods

2.1 Cox PH models

The notation used for Cox PH models (Cox, 1972) with one more subscript to capture multiple events is generalized. Let T_{ik} be the total time of the k^{th} event for the i^{th} subject, C_{ik} be the censoring time of the k^{th} event for the i^{th} subject. Let U_{ik} be the observation time, that is, $U_{ik} = \min(T_{ik}; C_{ik})$, and $\delta_{ik} = I(T_{ik} \leq C_{ik})$ is an indicator of observed k^{th} failure time for subject i . $Z_{ik} = (Z_{1ik}, Z_{2ik}, \dots, Z_{pik})$ is the covariate vector for the i^{th} subject with respect to the k^{th} event, and $Z_i = (Z'_{i1}, Z'_{i2}, \dots, Z'_{ik})$ denotes the covariate vector for the i^{th} subject, where K is the

maximum number of events within a subject. $\beta = (\beta_1, \beta_2, \dots, \beta_p)$ is a $p \times 1$ vector of unknown parameters. Denote $h_k(t|Z_i(t))$ as the hazard function for the k^{th} event of the i^{th} subject at time t . This is in the context of competing risk. In general, the hazard function at time t for a subject is defined as the instantaneous probability of failure at time t given the survivorship prior to time t and the covariates:

$$h_{k+1}(t | Z_i(t)) = \lim_{\Delta t \rightarrow 0} \Pr(t \leq T_{i,k+1} < t + \Delta t | T_{i,k+1} \geq t, Z_i(t)) / \Delta t. \quad (1)$$

Note that Cox PH model for the k^{th} event time T_k is

$$h_k(t|Z_i(t)) = h_{0,k}(t) \exp\{\beta' Z_i(t)\} \quad (2)$$

2.2 Correlated Gamma Frailty (CGF) Model

This model was introduced by Yashin and Iachine (1995a,b, 1997, 1999a,b) and applied to related lifetimes in many different settings. Examples are found in Pickles et al. (1994), Yashin et al. (1996), Iachine et al. (1998), Iachine (2002), Petersen (1998), Wienke et al. (2000, 2001, 2002, 2003a,b, 2004, 2005), Zdravkovic et al. (2002, 2004).

Let k_0, k_1, k_2 be some real positive values. Set $\lambda_1 = k_0 + k_1$ and $\lambda_2 = k_0 + k_2$

Let Y_0, Y_1, Y_2 be independently gamma distributed random variables with

$$Y_0 \sim \Gamma(k_0, \lambda_0), \quad Y_1 \sim \Gamma(k_1, \lambda_1), \quad Y_2 \sim \Gamma(k_2, \lambda_2) \quad \dots \quad \dots \quad \dots \quad (3)$$

Consequently,

$$Z_1 = \frac{\lambda_0}{\lambda_1} Y_0 + Y_1 \sim \Gamma(k_0 + k_1, \lambda_1) \quad \dots \quad \dots \quad \dots \quad (4)$$

$$Z_2 = \frac{\lambda_0}{\lambda_2} Y_0 + Y_2 \sim \Gamma(k_0 + k_2, \lambda_2) \quad \dots \quad \dots \quad \dots \quad (5)$$

$$\text{and } E(Z_1) = E(Z_2) = 1,$$

$$V(Z_1) = \frac{1}{\lambda_1} := \sigma_1^2, \quad (6)$$

$$V(Z_2) = \frac{1}{\lambda_2} := \sigma_2^2.$$

The following relation holds

$$\begin{aligned} E(Y_0^2) &= V(Y_0) + (E(Y_0))^2 \\ &= \frac{k_0}{\lambda_0^2} + \left(\frac{k_0}{\lambda_0}\right)^2 \quad \dots \quad \dots \quad (7) \\ &= \frac{k_0^2 + k_0}{\lambda_0^2} \end{aligned}$$

$$\begin{aligned}
 E(Z_1 Z_2) &= E\left(\frac{\lambda_0}{\lambda_1} Y_0 + Y_1\right) \left(\frac{\lambda_0}{\lambda_2} Y_0 + Y_2\right) \\
 &= E\left(\frac{\lambda_0^2}{\lambda_1 \lambda_2} Y_0^2 + \frac{\lambda_0}{\lambda_1} Y_0 Y_2 + \frac{\lambda_0}{\lambda_2} Y_0 Y_1 + Y_1 Y_2\right) \\
 &= \frac{\lambda_0^2}{\lambda_1 \lambda_2} \frac{k_0^2 + k_0}{\lambda_0^2} + \frac{\lambda_0 k_0 k_2}{\lambda_1 \lambda_0 \lambda_2} + \frac{\lambda_0 k_0 k_1}{\lambda_2 \lambda_0 \lambda_1} + \frac{k_1 k_2}{\lambda_1 \lambda_2} \quad (8) \\
 &= \frac{k_0 + (k_0 + k_1)(k_0 + k_2)}{\lambda_1 \lambda_2} \\
 &= \frac{k_0}{(k_0 + k_1)(k_0 + k_2)} + 1
 \end{aligned}$$

$$\begin{aligned}
 \text{Cov}(Z_1, Z_2) &= E(Z_1 Z_2) - E(Z_1)E(Z_2) \\
 &= \frac{k_0}{(k_0 + k_1)(k_0 + k_2)}, \quad \dots \dots \quad (9)
 \end{aligned}$$

This leads to the correlation

$$\begin{aligned}
 \rho &= \frac{\text{cov}(Z_1, Z_2)}{\sqrt{V(Z_1)V(Z_2)}} \quad \dots \dots \dots \quad (10) \\
 &= \frac{k_0}{\sqrt{(k_0 + k_1)(k_0 + k_2)}}
 \end{aligned}$$

Consequently, because of relation

$$\begin{aligned}
 k_0 + k_1 &= \lambda_1 \\
 &= 1/\sigma_i^2 \quad (i=1,2)
 \end{aligned}$$

It holds that

$$k_0 = \frac{\rho}{\sigma_1 \sigma_2} \quad \dots \dots \dots \quad (11)$$

and

$$\begin{aligned}
 k_i &= \frac{1}{\sigma_i^2} - k_0 \\
 &= \frac{1 - \frac{\sigma_i}{\sigma_j} \rho}{\sigma_i^2} \quad \dots \dots \dots \quad (12) \\
 & \quad (i, j = 1, 2; i \neq j).
 \end{aligned}$$

To derive the unconditional model, the Laplace transform of gamma distributed random variables is applied. Hence,

$$\begin{aligned}
 S(t_1, t_2) &= E\left\{S(t_1, t_2 | Z_1, Z_2)\right\} \\
 &= E\left\{S_1(t_1 | Z_1) S_2(t_2 | Z_2)\right\} \\
 &= E\left\{e^{-Z_1 \Lambda_1(t_1)} e^{-Z_2 \Lambda_2(t_2)}\right\} \\
 &= E\left\{e^{-\left(\frac{\lambda_0}{\lambda_1} Y_0 + Y_1\right) \Lambda_1(t_1)} e^{-\left(\frac{\lambda_0}{\lambda_2} Y_0 + Y_2\right) \Lambda_2(t_2)}\right\} \\
 &= E\left\{e^{-Y_0 \left(\frac{\lambda_0}{\lambda_1} \Lambda_1(t_1) + \frac{\lambda_0}{\lambda_2} \Lambda_2(t_2) - Y_1 \Lambda_1(t_1) - Y_2 \Lambda_2(t_2)\right)}\right\} \\
 &= \left(1 + \frac{1}{\lambda_0} \left(\frac{\lambda_0}{\lambda_1} \Lambda_1(t_1) + \frac{\lambda_0}{\lambda_2} \Lambda_2(t_2)\right)\right)^{-k_0} \left(1 + \frac{1}{\lambda_1} \Lambda_1(t_1)\right)^{-k_1} \left(1 + \frac{1}{\lambda_2} \Lambda_2(t_2)\right)^{-k_2} \cdot (13) \\
 &= \left(1 + \sigma_1^2 \Lambda_1(t_1) + \sigma_2^2 \Lambda_2(t_2)\right)^{-\frac{\rho}{\sigma_1 \sigma_2}} \left(1 + \sigma_1^2 \Lambda_1(t_1)\right)^{-\frac{1 + \frac{\sigma_1}{\sigma_2} \rho}{\sigma_1^2}} \left(1 + \sigma_2^2 \Lambda_2(t_2)\right)^{-\frac{1 + \frac{\sigma_2}{\sigma_1} \rho}{\sigma_2^2}}
 \end{aligned}$$

Which results in the representation of the Correlated Gamma Frailty model given as

$$S(t_1, t_2) = \frac{S_1(t_1)^{1 - \frac{\sigma_1}{\sigma_2} \rho} S_2(t_2)^{1 - \frac{\sigma_2}{\sigma_1} \rho} e^{\beta x_{ij}}}{\left(S_1(t_1) \sigma_1^2 + S_2(t_2) \sigma_2^2 - 1\right)^{\frac{\rho}{\sigma_1 \sigma_2}}}, \dots \dots \dots (14)$$

2.3 The Proposed Model – Modified Gamma Frailty (MGF) Model

In order to include heterogeneity in the model, we assume a correlated gamma frailty model. Let Z_j ($j = 1; 2$) be the frailties, and X_j ($j = 1; 2$) vectors of observable covariates of the two individuals of a twin pair. Assume that their individual hazards are represented by the proportional hazards model

$$\lambda(t) = Z_j \lambda_0(t) \exp\{\beta^T X_j\} \quad (j = 1, 2) \quad \dots \dots \dots (15)$$

with a baseline hazard function $\lambda_0(t)$ describing the risk of respiratory infection as a function of age and β denotes the vector of regression parameters. Let the lifetimes of the two twin partners be conditionally independent given their frailties Z_1 and Z_2 . Because frailties Z_j ($j = 1; 2$) are usually unobservable, their correlation coefficient used cannot be estimated directly from the empirical data. So a bivariate lifetime model which allows indirect calculation of the parameters is needed. The unconditional bivariate survival function of the correlated gamma frailty model with observed covariates is given by:

$$S(t_1, t_2 | X_1, X_2) = S(t_1 | X_1)^{1-\rho} S(t_2 | X_2)^{1-\rho} S(t_1 | X_1)^{-\sigma^2} + \{S(t_2 | X_2)^{-\sigma^2} - 1\} \frac{\rho}{\sigma^2} \dots \dots (16)$$

where $S(t | X)$ denotes the marginal univariate survival function, assumed to be equal for both partners in a twin pair. Using a parametric approach we fit a model to the data, such that

$$S(t | X_{ijk}) = \left[1 + \left[(1 + \sigma_1^2 \frac{a}{b} (e^{bt} - 1)) \frac{\sigma_2^2}{\sigma_1^2} \rho - 1 \right] \right]^{-\frac{1}{\sigma_1^2}} e^{\beta_k X_{ijk}} \dots \dots (17)$$

where $a, b, \sigma_1^2, \sigma_2^2, \beta$ and ρ are parameters to be estimated. The lifetimes are assumed to be independently censored from the right by independent and identically distributed pairs of non-negative random variables, which are independent of the lifetimes. Thus, observe

$$(T_{i1}, T_{i2}, \Delta_{i1}, \Delta_{i2}, X_{i1}, X_{i2}) \dots \dots \dots (18)$$

with Δ_{ij} ($i = 1, 2, \dots, n; j = 1, 2$) as a binary variable with values 1 (event) and 0 (no event). Let the lifetimes follow a distribution (dependent on covariates X_1, X_2) given by the bivariate survival function

$$S(t_1, t_2 | X_1, X_2) = P(T_{i1} > t_1, T_{i2} > t_2 | X_1, X_2) \dots \dots \dots (19)$$

Starting from this model, we are able to derive the likelihood function given by

$$L(t_1, t_2, \delta_1, \delta_2, X_1, X_2) = \delta_1 \delta_2 S_{t_1 t_2}(t_1, t_2 | X_1, X_2) - \delta_1 (1 - \delta_2) S_{t_1}(t_1, t_2 | X_1, X_2) - (1 - \delta_1) \delta_2 S_{t_2}(t_1, t_2 | X_1, X_2) + (1 - \delta_1)(1 - \delta_2) S(t_1, t_2 | X_1, X_2) \dots \dots (20)$$

Partial derivatives of the marginal survival functions are given by

$$S_{t_j}(t_1, t_2) = \frac{\partial S(t_1, t_2)}{\partial t_j} \quad (j = 1, 2) \dots \dots \dots (21)$$

and

$$S_{t_1, t_2}(t_1, t_2) = \frac{\partial S(t_1, t_2)}{\partial t_1 \partial t_2} \quad (22)$$

The model is called the **Modified Gamma Frailty (MGF) Model**.

3. Numerical Illustration

An application of the models to an existing data on occupational exposure tagged – IRANIAN data is demonstrated here. Relationships between occupational exposures and morbidity, morbidity and job category were analyzed using proportional hazard analysis, allowing for exposure status (never exposed, ever smoked and ever exposed) until the time of carrying out the study. The survival-analysis was performed using the SPSS VERSION 15. The discrete algorithm was used, since the time-scale (person-years) was discrete. All exposures were first analyzed separately, allowing for age and smoking habits. Two-sided p-values < 0.05 were considered as statistically significant. The relationship between occupational exposures and morbidity was also analyzed simultaneously. Using the stepwise option of SPSS, and allowing for age and smoking habits, specific exposures were included and excluded until the following conditions were met: the significance of the residual Chi-squared was less than 0.25, and the significance of the relative risks was less than 0.10. Using the standard error of the regression coefficient, the 95% confidence intervals were estimated. The Matlab software and R was also applied in analyzing the Correlated Gamma Frailty Model and the Modified Gamma Frailty Model. Hazard function and survival functions for the exposure data for large and small samples were estimated.

4. Results

Tables 1-4 shows the results of analysis of the Iranian data and the goodness of fit table. The exponentiated coefficients in the third column of each table of the output shown are interpretable as multiplicative effects on the hazard. In tables 1, for example, holding the other covariates constant, one additional year of age increases the yearly hazard of exposure of worker by a factor of $e^\beta = 1.053376$ on average – that is, by 5.3 percent. Similarly, each Forced Ventilatory Function (FVC) factor increases the hazard by a factor of 1.059079 or 5.9 percent. The fifth column is the result of the test of significance of β using the Wald Statistic which is the ratio of the coefficients to the standard error of β . The obtained value is compared with the Z value and a decision is made. In tables 2, holding the other covariates constant, an additional year of age increases the yearly hazard of exposure of worker by a factor of $e^\beta = 1.034585$ on average – that is, by 3.5 percent. Similarly, each FVC factor increases the hazard by a factor of 1.001301 or 0.1 percent. In tables 3, holding the other covariates constant, an additional year of age increases the yearly hazard of exposure of worker by a factor of $e^\beta = 1.053481$ on average – that is, by 5.3 percent. Similarly, each FVC factor increases the hazard by a factor of 1.062155 or 6.2 percent. The exposure status (never exposed, exposed and ever smoked), Job category and pack years smoked is considered to be insignificant for the Iranian data using the Cox Model. The CGF

captures the exposure status and Job category to be insignificant for the Iranian data while the proposed MGF considers all the variables to be significant for the Iranian data.

Covariate	coef $f(\beta)$	Exp(coef $f(\beta)$)	Std error Coef $f(\beta)$	Z	P	95% C.I for coef $f(\beta)$
AGE	0.0520	1.053376	0.0227	2.290749	0.0010	1.0212- 1.0828
BMI	0.0555	1.057069	0.0119	4.663866	0.0001	1.0321- 1.0788
EXPOSURE STATUS	- 0.1498	0.86088	0.2122	-0.70594	0.2145*	0.1531 - 1.9214
JOB CATEGORY	0.4337	1.542956	0.3819	1.135638	0.1041*	0.2009 - 1.6029
SYST B P	0.0915	1.095817	0.0286	3.199301	0.0029	1.0899 - 1.1103
PACK YRS SMOKED	- 0.2038	0.815625	0.1914	-1.06479	0.9856*	0.2133- 1.9339
FVC	0.0574	1.059079	0.0221	2.597285	0.0085	1.0444 - 1.0679
FEV ₁	0.0849	1.088608	0.1956	0.434049	0.0007	1.0415 - 1.0940

* Not significant.

Tables 1: Regression Coefficients in the Cox Model for the Iranian Study

Covariate	coef $f(\beta)$	Exp(coef $f(\beta)$)	Std error coef $f(\beta)$	Z	P	95% C.I for coef $f(\beta)$
AGE	0.0340	1.034585	0.0154	2.207792	0.0011	1.0112- 1.0542
BMI	0.0437	1.044669	0.0231	1.891775	0.0053	1.0221 -1.0658
EXPOSURE STATUS	- 0.0249	0.975407	0.0713	-0.34923	0.6714*	0.0131 -1.0224
JOB CATEGORY	0.0023	1.002303	0.0044	0.522727	0.1304*	0.0019-1.0329
SYST B P	0.0021	1.002102	0.0009	2.333333	0.0039	1.0001 -1.0030
PACK YRS SMOKED	- 0.2138	0.80751	0.0371	-5.7628	0.0035	0.033 - 0.9339
FVC	0.0013	1.001301	0.0006	2.16667	0.0018	1.0003- 1.0489
FEV ₁	0.0362	1.036863	0.0146	2.479452	0.0029	1.0235 -1.0440

* Not significant.

Tables 2: Regression Coefficients in the Correlated Gamma Frailty Model for the Iranian Study

Covariate	coef $f(\beta)$	Exp(coef $f(\beta)$)	Std error coef $f(\beta)$	Z	P	95% C.I for coef $f(\beta)$
AGE	0.0521	1.053481	0.0167	3.11976	0.0011	1.0215 - 1.0848
BMI	0.0547	1.056224	0.0111	4.927928	0.0001	1.0324 - 1.0798
EXPOSURE STATUS	- 0.1049	0.900415	0.2013	-0.52111	0.0045	0.0312 -0.9814
JOB CATEGORY	0.3987	1.489887	0.4009	0.994512	0.0041	1.0009-1.6029
SYST B P	0.0891	1.09319	0.0271	3.287823	0.0021	1.0021 -1.5003
PACK YRS SMOKED	- 0.2038	0.815625	0.1717	-1.18695	0.0056	0.633 - 0.9939
FVC	0.0603	1.062155	0.0211	2.85782	0.0015	1.0044 - 1.2479
FEV ₁	0.0762	1.079178	0.0377	2.02122	0.0029	1.0135 -1.8340

* Not significant.

Tables 3: Regression Coefficients in the Modified Gamma Frailty Model for the Iranian Study

Prognostic factors	Cox regression HR (CI 95%)	Correlated Gamma Frailty HR (CI 95%)	Modified Gamma Frailty HR† (CI§ 95%)
Age	1.0530 (1.0212 - 1.0828)	1.0346 (1.0112 - 1.0542)	1.053481(1.0215 - 1.0848)
BMI	1.0571(1.0321 - 1.0788)	1.0447(1.0221 -1.0658)	1.056224(1.0324 - 1.0798)
EXPOSURE STATUS	0.8609(0.1531 - 1.9214)*	0.9754 (0.0131 -1.0224)*	0.900415(0.0312 -0.9814)
JOB CATEGORY	1.5430 (0.2009 - 1.6029)*	1.0023 (0.0019-1.0329)*	1.48989(1.0009-1.6029)
SYST B P	1.0958 (1.0899 - 1.1103)	1.0021 (1.0001 -1.0030)	1.0932(1.0021 -1.5003)
PACK YRS SMOKED	0.8156 (0.2133 - 1.9339)*	0.8075 (0.033 - 0.9339)	0.8156(0.633 - 0.9939)
FVC	1.0591(1.0444 -1.0679)	1.0013 (1.0003 - 1.0489)	1.062155(1.0044 - 1.2479)
FEV ₁	1.0886 (1.0415 - 1.0940)	1.0369(1.0235 -1.0440)	1.079178(1.0135 - 1.8340)
AIC#	1,157	751	704

† Hazard Ratio § Confidence interval * Not significant # Akaike Information Criterion
 Tables 4 : Prognostic Factors of Occupational Exposure using Cox and frailty Models for Iranian study




Fig. 1 Survival Function at mean of covariates - Iranian Study




Fig. 2 Hazard Function at mean covariates - Iranian Study

5. Conclusion

Interestingly, parameter estimates are quite different depending on distribution of the base-line hazard function. The newly introduced Modified Gamma frailty model offers a very elegant approach to integrate the concept of clusters into frailty modelling. The survival function is explicitly available and of easy form which allows traditional maximum likelihood parameter estimation. This is the most important advantage of the suggested model compared to the model introduced by Moger and Aalen (2005). The present work contributes to three aspects of Frailty

models with censored data. First, It presents several important extensions of the existing models. Secondly, It develops a general asymptotic theory for the Frailty models. Thirdly, It provides simple and efficient numerical method to implement the corresponding inference procedures. It is hoped that this work will facilitate further development and applications of Frailty models. It has been demonstrated that the MGF is a very general and powerful approach to the analysis of Frailty models with censored data. This approach can be used to study many other problems. Of great interest would be a non-parametric version of the correlated compound Poisson frailty model, where the baseline hazard functions are not specified. A part of future research is envisaged in this direction. Another aspect that will be of interest for further research is the problem of identifiability. The identifiability problem is growing with increased censoring, but is reduced by the parametric modelling of the baseline hazard.

References

- Aalen, O. O. (1988). Heterogeneity in survival analysis. *Statistics in Medicine*, 7, 1121- 1137.
- Aalen, O. O. (1992). Modelling heterogeneity in survival analysis by the compound poisson distribution. *Annals of Applied Probability*, 4, 951-972.
- Anthony I. W., Isaac D. E., Victor A. K., (2019): Survival Analysis of Under –five Mortality and Its Associated Determinants in Nigeria: Evidence from a Survey Data *International Journal of Statistics and Applications* 9(2): 59-66.
- Cox, D.R. (1972). Regression models and life tables, *Journal of the Royal Statistical Society, B* 34, 187-400.
- Cox, D.R. (1975). Partial likelihood. *Biometrika*, 62, 269-276.
- Getachew Y. & Bekele S., (2016). Survival Analysis of Under-Five Mortality of Children and its Associated Risk Factors in Ethiopia. *J Biosens Bioelectron* 7: 213.
- Iachine, I., Holm, N., Harris, J., Begun, A., Iachina, M., Laitinen, M., Kaprio, J., Yashin, A. (1998). How heritable is individual susceptibility to death? The results of an analysis of survival data on Danish, Swedish and Finnish twins. *Twin Research* 1, 196 – 205.
- Iachine, I. (2002). The Use of Twin and Family Survival Data in the Population Studies of Aging: Statistical Methods Based on Multivariate Survival Models. Ph.D. Thesis. Monograph 8, Department of Statistics and Demography, University of Southern Denmark.
- Kalbfleisch, J. D. and R.L. Prentice, (1980). The statistical analysis of failure time data (John Wiley & Sons Inc., New York).
- Kiefer, N.M., (1988). Economic duration data and hazard functions, *Journal of Economic Literature* 26, 646-679.
- Lancaster, T., (1990). The econometric analysis of transit data (Cambridge University Press, Cambridge, U.K).
- Lee T.E and Wang J. W., (2003). *Statistical Methods for Survival Data Analysis*. 3rd ed. A John Wiley and Sons, Inc., Publication.
- Longini, I. M., and Halloran, M. E. (1996). A frailty mixture model for estimating vaccine efficacy. *Applied Statistics*, 45, 165-173.
- Moger, T. A., and Aalen, O. O. (2005). A distribution for multivariate frailty based on the compound poisson distribution with random scale. *Lifetime Data Analysis*, 11, 41-59.
- Petersen J. H (1998). An Additive Frailty Model for Correlated Life Times," *Biometrics* 54 : 646-661.

- Petersen, T., (1991). Time aggregation bias in continuous-time hazard-rate models.
- Pickles, A., Crouchley, R., Simono, E., Eaves, L., Meyer, J., Rutter, M., Hewitt, J., Silberg, J. (1994). Survival models for developmental genetic data: age of onset of puberty and antisocial behavior in twins. *Genetic Epidemiology* 11, 155 – 170.
- Price, D. L., and Manatunga, A. K. (2001). Modelling survival data with a cured fraction using frailty models. *Statistics in Medicine*, 20, 1515-1527.
- Vermunt, J.K., (1996). Log-linear event history analysis (Tilburg University Press, Netherlands).
- Vilcassim, N. J. and D.C. Jain, (1991). Modeling purchase timing and brand switching behavior incorporating explanatory variables and unobserved heterogeneity, *Journal of Marketing Research*, 28, 29-41.
- Wedel, M., W.A. Kamakura, W.S. DeSarbo and F. ter Hofstede, (1995). Implications for asymmetry, non proportionality and heterogeneity in brand switching models from piece-wise exponential mixture hazard models, *Journal of Marketing Research*, 32, 457-463.
- Wienke, A., Christensen, K., Holm, N., Yashin, A. (2000). Heritability of death from respiratory diseases: an analysis of Danish twin survival data using a correlated frailty model. Medical Infobahn for Europe. A. Hasman et al., IOS Press, Amsterdam, 407-411.
- Wienke, A., Holm, N., Skytthe, A., Yashin, A.I. (2001). The heritability of mortality due to heart diseases: a correlated frailty model applied to Danish twins. *Twin Research* 4, 266 – 274.
- Wienke, A., Christensen, K., Skytthe, A., Yashin, A.I. (2002). Genetic analysis of cause of death in a mixture model with bivariate lifetime data. *Statistical Modelling* 2, 89 – 102.
- Wienke, A., Lichtenstein, P., Yashin, A.I. (2003a). A bivariate frailty model with a cure fraction for modeling familial correlations in diseases. *Biometrics* 59, 1178 – 1183.
- Wienke, A., Holm, N., Christensen, K., Skytthe, A., Vaupel, J., Yashin, A.I. (2003b). The heritability of cause-specific mortality: a correlated gamma-frailty model applied to mortality due to respiratory diseases in Danish twins born 1870 - 1930. *Statistics in Medicine* 22, 3873 – 3887.
- Wienke, A. (2004). Die Vererbbarkeit der Todesursache: ein correlated-frailty Modell angewendet auf Danische Zwillinge, geboren 1870 - 1930. In: R. Scholz und J. Flathmann (Hrsg): Lebenserwartung und Mortalität. Materialien zur Bevölkerungs- und Gesundheitswissenschaft Heft 111, Wiesbaden, 81-98.
- Yashin, A.I., Iachine, I.A. (1995a). Genetic analysis of durations: Correlated frailty model applied to survival of Danish twins. *Genetic Epidemiology* 12, 529 – 538.
- Yashin, A.I., Iachine, I.A. (1995b). Survival of related individuals: an extension of some fundamental results of heterogeneity analysis. *Mathematical Population Studies* 5, 321-39.
- Yashin, A.I., Manton, K.G., Iachine, I.A. (1996). Genetic and environmental factors in duration studies: multivariate frailty models and estimation strategies. *Journal of Epidemiology and Biostatistics* 1, 115 – 120.
- Yashin, A.I., Iachine, I.A. (1997). How frailty models can be used for evaluating longevity limits: Taking advantage of an interdisciplinary approach. *Demography* 34, 31 – 48.
- Yashin, A.I., Iachine, I. (1999a). Dependent hazards in multivariate survival problems. *Journal of Multivariate Analysis* 71, 241 – 261.
- Yashin, A.I., Iachine, I. (1999b). What difference does the dependence between durations make? Insights for population studies of aging. *Lifetime Data Analysis* 5, 5 – 22.
- Wienke, A., Arbeev, K., Locatelli, I., Yashin, A.I. (2005). A comparison of different correlated frailty models and estimation strategies. *Mathematical Biosciences* 198, 1 – 13.

Zdravkovic, S., Wienke, A., Pedersen, N.L., Marenberg, M.E., Yashin, A.I., de Faire, U. (2002). Heritability of death from coronary heart disease: a 36 years follow-up of 20,966 Swedish twins. *Journal of Internal Medicine* 252, 247 – 54.

Zdravkovic, S., Wienke, A., Pedersen, N.L., Marenberg, M.E., Yashin, A.I., de Faire, U. (2004). Genetic influences on CHD-death and the impact of known risk factors: Comparison of two frailty models. *Behavior Genetics* 34, 585 – 591.